JUNIOR ICE HOCKEY · PHYSICAL PREPARATION

Want to win more games?

5 things nearly every ice hockey program does not coach.

Prepared by John Bryan & Candace Nakagawa
Empyrean Health & Wellness, Soldotna, AK
Email: Team@EmpyreanFitness.com · Website: EmpyreanFitness.com

Most programs train the wrong things — or the right things, wrong.

Most junior hockey programs are training the wrong things, or training the right things with the wrong methods.

The 16–21 age range is arguably the highest-leverage period for neuromuscular and aerobic development in an athlete's career — and it's routinely underutilized because game schedules dominate at the expense of systematic physical preparation.

IN THIS REPORT

1. Neuromuscular Power Output & Rate of Force Development
2. Alactic-Anaerobic Energy System Capacity & Resynthesis Rate
3. Hip & Groin Force Production in the Frontal Plane
4. Ventilatory Threshold & Aerobic Base Quality
5. Structural Readiness & Tissue Tolerance

01 Neuromuscular Power Output & Rate of Force Development (RFD)

This is the single biggest limiter we see in this age group. At 16–21, the nervous system is still maturing — specifically the myelination of fast-twitch motor pathways. Players have muscle mass but can't yet recruit it explosively on demand.

WHY IT LIMITS PERFORMANCE

- First-step acceleration (0–5m) is almost entirely a neuromuscular event, not a metabolic one
- Checking battles and board play require instantaneous force expression that immature motor patterns can't produce
- Most junior programs still train for hypertrophy rather than rate of force development

WHAT THE SCIENCE SHOWS

Elite players demonstrate significantly faster RFD profiles than age-matched juniors — the gap is largely neurological, not structural. Triphasic and complex/contrast methods are chronically underutilized at this level.

HOW TO FIX IT

Contrast / complex training
Pair a heavy strength movement (trap-bar deadlift, back squat) with an explosive movement (broad jump, med ball throw), 2x/week.

Velocity-based training (VBT)
Track bar speed on key lifts so athletes train explosive intent at lighter loads (30–60% 1RM), not just heavy grinding reps.

Reactive / plyometric work
Add depth jumps, skater bounds, and short (0–5m) resisted or unresisted sprint starts twice weekly.

Separate hypertrophy & RFD blocks
Sequence 4–6 week hypertrophy phases followed by 3–4 week power/RFD phases so strength converts into speed.

02 Alactic-Anaerobic Energy System Capacity & Resynthesis Rate

Hockey shifts are 35–50 seconds of repeated maximal or near-maximal bursts. The phosphocreatine (PCr) system drives those efforts — but PCr resynthesis between shifts is where most junior players fall apart.

WHY IT LIMITS PERFORMANCE

- PCr resynthesis is ~50% complete in 30 seconds and ~95% in ~3 minutes — bench rest is often only 60–90 seconds
- Players who can't rapidly resynthesize PCr see a 15–25% drop in peak power by the 2nd period
- The aerobic system feeds PCr resynthesis — your aerobic ceiling governs your top-end explosive repeatability

WHAT THE SCIENCE SHOWS

This is the "aerobic-anaerobic interface" — most teams train one or the other, not the system that bridges them. Repeated sprint ability (RSA) training is severely underrepresented in junior programs.

HOW TO FIX IT

Match conditioning to shift demands
Build repeated-sprint intervals using hockey work:rest ratios (30–40s max effort, 60–90s recovery), not steady-state cardio.

Train the 'interval within the interval'
Sets of 3–5 short maximal efforts (5–10s) with brief 20–30s rests inside a longer recovery block, mimicking bench-to-shift demand.

Build the aerobic engine first
A higher aerobic base speeds PCr resynthesis — layer in tempo/aerobic work before adding anaerobic volume.

Track fatigue drop-off
Use repeat-sprint tests (6x 30m) and watch %decrement across reps, not just the fastest single rep.

03 Hip & Groin Force Production in the Frontal Plane

The most sport-specific structural limiter that nobody talks about enough. Skating power is generated through hip abduction and adduction in the frontal plane — not the sagittal plane most strength programs are built around.

WHY IT LIMITS PERFORMANCE

- Adductor-to-abductor strength ratios are frequently imbalanced, robbing stride power and raising groin injury risk
- Skating stride is a lateral push — hip external rotator and abductor weakness leaks energy at the hip every stride
- Pubescent growth spurts alter hip geometry rapidly, disrupting motor patterns mid-season

WHAT THE SCIENCE SHOWS

Copenhagen adduction exercises, lateral band work, and single-leg frontal plane loading are not standard in most junior programs. Players are squatting but not skating-pattern loading.

HOW TO FIX IT

Copenhagen adduction progressions
2x/week, from bent-knee/short-lever to long-lever, to close the adductor-to-abductor gap and cut groin injury risk.

Lateral & rotational strength work
Lateral band walks, skater bounds, cossack squats, single-leg lateral step-ups train the frontal-plane pattern.

Copy the skating pattern under load
Resisted lateral sled pushes or slideboard work reinforce hip abduction/adduction in the exact plane the ice demands.

Screen through growth spurts
Periodic hip strength/ROM screening for 16–18 year-olds catches asymmetries early during rapid growth phases.

04 Ventilatory Threshold & Aerobic Base Quality

VO2max gets all the attention, but ventilatory threshold (VT2) is the real performance predictor — the highest sustainable work rate before blood lactate accumulates uncontrollably.

WHY IT LIMITS PERFORMANCE

- Low VT2 players accumulate lactate faster, perceive effort higher, and make worse decisions under fatigue
- Aerobic base in 16–18 year olds is chronically underdeveloped — youth hockey prioritizes skill over aerobic conditioning
- Off-ice aerobic training is inconsistently programmed — players arrive with wildly variable aerobic capacities

WHAT THE SCIENCE SHOWS

The best NHL prospects often test with off-ice VO2max of 60–65+ ml/kg/min. The average junior arriving at a USHL/OHL camp is often 52–57 — months of aerobic base work that wasn't done.

HOW TO FIX IT

Build a real off-season aerobic base
2–3 sessions/week of steady Zone 2 work (bike, run, row) for 6–8 weeks before ramping anaerobic/on-ice volume.

Add threshold-specific tempo intervals
Once a base exists, use longer tempo efforts (4x8min at 'comfortably hard' pace) to shift VT2 upward.

Test and track, don't guess
A field test (20–30min time trial or step test with HR/RPE) each off-season gives a baseline to track against.

Protect the aerobic block
Schedule aerobic sessions early in the off-season so they aren't the first thing cut when games ramp up.

05 Structural Readiness & Tissue Tolerance (Tendon / Bone Maturity)

The most underappreciated limiter in this age group — invisible until it becomes an injury.

WHY IT LIMITS PERFORMANCE

- Tendons and bones adapt more slowly than muscle — connective tissue lags 6–18 months behind puberty-driven strength gains
- The growth plate at the apophysis is biologically weaker than the surrounding tendon — Osgood-Schlatter and Sever's are epidemic
- Players cannot train needed volumes because tissue tolerance, not motivation or muscle, is the rate-limiter

WHAT THE SCIENCE SHOWS

This is a biological debt situation. Aggressive early specialization starves bone and tendon of varied loading, so players arrive with structurally immature tissue that breaks down under the step-up in training and game load.

HOW TO FIX IT

Ramp training load gradually
Cap week-to-week volume/intensity increases (~10% max) to give tendon and bone time to catch up to muscle.

Add tendon-specific loading
Slow heavy-resistance isometrics and eccentrics (calf raises, seated knee extensions) build tendon stiffness, not just muscle size.

Push back on early over-specialization
Encourage multi-sport or cross-training so bone and connective tissue get diverse loading stimulus.

Watch growth-spurt windows
Monitor for apophyseal pain (Osgood-Schlatter, Sever's), back off volume proactively, support recovery with protein, calcium, vitamin D.

The five limiters, at a glance

LIMITER	PRIMARY SYSTEM	TRAINABILITY, 16–21	TYPICAL NEGLECT
Neuromuscular RFD	CNS / Motor	High	Very High
PCr Resynthesis Rate	Alactic-Aerobic Interface	High	High
Frontal Plane Hip Force	Musculoskeletal	High	Very High
Ventilatory Threshold	Cardiovascular	Very High	Moderate
Tissue Tolerance	Connective Tissue	Moderate	Very High

Sequence contrast training, repeated-sprint conditioning, frontal-plane hip work, aerobic base building, and gradual tissue loading — in that order of urgency — across the off-season calendar.

END OF REPORT